

*Полтавський державний медичний університет
Полтавське відділення Міжнародного фонду допомоги хворим
з наслідками травм та захворювань
Всеукраїнська громадська організація «Наукове товариство анатомів,
гістологів, ембріологів та топографоанатомів України»*

DOI 10.26724
ISSN 2079-8334
E-ISSN 2412-9348

Світ медицини та біології

№ 2 (80) 2022

Науковий, медичний, екологічний журнал

**Заснований в травні 2005 року
Виходить 4 рази на рік**

Полтава • 2022

Засновник
Полтавський державний медичний
університет

Свідоцтво про державну реєстрацію
КВ № 9878 від 23.05.2005

Фахове наукове видання України
(Наказ МОН України № 612 від 07.05.2019 р.)
Медичні і біологічні науки

Чайковський Ю.Б. (Київ) – головний редактор
Ждан В.М. (Полтава) – заступник головного редактора
Шепітько В.І. (Полтава) – заступник головного редактора
Єрошенко Г.А. (Полтава) – відповідальний редактор

Редакційна колегія:

Аветіков Д.С. (Полтава), **Борнштейн Натан** (Тель-Авів), **Валіуліс Арунас** (Вільнюс),
Вастьянов Р.С. (Одеса), **Геращенко С.Б.** (Івано-Франківськ), **Голованова І.А.** (Полтава),
Громова А.М. (Полтава), **Дворник В.М.** (Полтава), **Костенко В.О.** (Полтава),
Костиленко Ю.П. (Полтава), **Крючко Т.О.** (Полтава), **Ліхачов В.К.** (Полтава), **Ляховський В.І.** (Полтава), **Мишковска Дорота** (Ягеллонськ), **Наркевич Кжиштоф** (Гданськ),
Похилько В.І. (Полтава), **Родінкова В.В.** (Вінниця), **Сілкіна Ю.В.** (Дніпро), **Скрипник І.М.** (Полтава),
Скрипніков А.М. (Полтава), **Сокуренко Л.М.** (Київ),
Старченко І.І. (Полтава), **Ткаченко П.І.** (Полтава), **Фал Анджей Маріуш** (Варшава),
Шерстюк О.О. (Полтава)

Рекомендовано Вченою радою ПДМУ (протокол № 6 від 9.03.2022 р.)

Відповідальний за випуск – Єрошенко Г.А.
Комп'ютерна верстка – Наріжна О.М.
Наукове редактування – редакція

Включений до науково-метричної бази даних **WEB OF SCIENCE**
Розміщений на онлайнових базах даних **PROQUEST, INDEX COPERNICUS**
та **GOOGLE SCHOLAR**

Адреса редакції та видавця –
Полтавський державний медичний університет,
кафедра гістології, цитології та ембріології,
вул. Шевченка, 23, м. Полтава, 36000
Тел. (05322) 60-84-44. E-mail: womab.ed@gmail.com

Сайт журналу – www.womab.com.ua

CONTENTS

CLINICAL MEDICINE

Azimov E.H.

Comparative analysis of circumferential resection margin condition during the laparoscopic and open total mesorectal excision

Atamaniuk O.Yu., Skrypko V.D.,

Atamaniuk V.M., Bybliuk Yu.Y.

Relationship between ultrasonographic criteria, clinical manifestations and life quality parameters in patients with lower extremity varicose vein disease

Balytsky V.V., Zakharash M.P., Kuryk O.G.

Clinical and morphological evaluation of the effectiveness of using radio wave and high-frequency electrosurgery technologies for the treatment of combined anorectal diseases

Bilovol O.M., Kniazkova I.I., Kirienko O.M.,

Kuzminova N.V., Golovachova V.O.,

Kirienko D.O., Abramova L.P.

Functional condition of kidneys in patients with arterial hypertension in combination with type 2 diabetes mellitus

Bukach O.P., Buzdugan I.O.

Dyslipidemia in patients with rheumatoid arthritis depends on comorbid pathology and genetic predictors

Varzhapetian S.D., Kovach I.V., Sydor O.V.,

Stroganova T.V., Buniatian K.A., Dats V.V.,

Kucherenko A.N.

Severity of adentia as a risk factor of repeated dental implant operations

Hajiyeva Y.H.

Comparative evaluation of the effectiveness of modern means of hypercholesterolemia correction

Horodova-Andrieieva T.V., Liakhovskyi V.I.,

Sydorenko A.V.

Influence of vacuum instillations with intravenous introduction of nitrogen oxide donators on the healing of purulent soft tissue wounds

Horokhovskyi V.V., Dienha O.V., Dienga A.E.,

Pyndus T.O., Jenča A. Jr., Shnaider S.A.,

Tsushko I.O.

Analysis of the frequency and structure of non-carious lesions of hard tissues of teeth in children

Hrytsuliak B.V., Hrytsuliak V.B., Bielova N.V.,

Hlodan O.Ia., Ivasiuk I.Y., Mykytyn T.V.

Structural and functional changes in the prostate gland in diabetes mellitus

Dvornyk A.V., Dvornyk V.M., Vodoriz Y.Y.,

Skripnykov P.M., Tkachenko I.M.

Clinical features of influence of different groups of bleaching agents in the oral cavity at different periods after the end of the treatment

Dienha O.V., Malekh N.V., Makarenko O.A.,

Pyndus T.O., Jenča A., Shnaider S.A.

Biochemical parameters of oral fluid of children with hypertrophic gingivitis during dental treatment

Zak M.Y., Kiro L.S., Kushnirenko I.V.,

Yablonska T.M., Airapetian I.A., Sverdlova M.V.

Peculiarities of the course of somatized and anxiety disorders in patients with non-alcoholic fatty liver disease on the background of abdominal obesity

ЗМІСТ

Клінічна медицина

- 7 Азімов Е.Г.
Порівняльний аналіз стану краю кругової резекції при лапароскопічній і відкритій тотальній мезоректальній ексцизії
- 13 Атаманюк О.Ю., Скрипко В.Д.,
Атаманюк В.М., Библиук Ю.Й.
Взаємозв'язок ультрасонографічних критеріїв, клінічних проявів та параметрів якості життя у пацієнтів із варикозною хворобою нижніх кінцівок
- 19 Балицький В.В., Захараш М.П., Курик О.Г.
Клінічна та морфологічна оцінка ефективності використання сучасних радіохвильових та високочастотних електрохірургічних технологій для лікування комбінованих аноректальних захворювань
- 24 Біловол О.М., Князькова І.І., Кіріченко О.М.,
Кузьмінова Н.В., Головачова В.О.,
Кіріченко Д.О., Абрамова Л.П.
Функціональний стан нирок у пацієнтів з артеріальною гіпертензією у поєднанні з цукровим діабетом 2 типу
- 29 Букач О.П., Буздуган І.О.
Дисліпідемія у хворих на ревматоїдний артрит залежно від коморбідної патології та генетичних предикторів
- 33 Варжапетян С.Д., Ковач І.В., Сидор О.В.,
Строгонова Т.В., Бунятян К.А., Дац В.В.,
Кучеренко О.М.
Протяжність дефекту зубного ряду як фактор ризику повторної операції з дентальної імплантації
- 38 Гаджиєва Я.Г.
Порівняльна оцінка ефективності сучасних засобів корекції гіперхолестеринемії
- 43 Городова-Андрієва Т.В., Ляховський В.І.,
Сидоренко А.В.
Вплив вакуумістиляції з внутрішньовенным введенням донаторів оксиду азоту на загоєння гнійних ран м'яких тканин
- 47 Гороховський В.В., Деньга О.В., Деньга А.Е.,
Піндус Т.О., Єнча А., Шнайдер С.А.,
Цушко І.О.
Аналіз частоти та структури некаріозних уражень твердих тканин зубів у дітей
- 51 Грицуляк Б.В., Грицуляк В.Б., Белова Н.В.,
Глодан О.Я., Івасюк І.Й., Мікитин Т.В.
Структурно-функціональні зміни в передміхуровій залозі при цукровому діабеті
- 56 Дворник А.В., Дворник В.М., Водоріз Я.Ю.,
Скрипников П.М., Ткаченко І.М.
Клінічні особливості впливу вибілювачів різних груп у порожнині рота у різні терміни після закінчення лікування
- 60 Деньга О.В., Малех Н.В., Макаренко О.А.,
Піндус Т.О., Єнча А., Шнайдер С.А.
Біохімічні показники ротової рідини дітей з гіпертрофічним гінгівітом у процесі стоматологічного лікування
- 65 Зак М.Ю., Кіро Л.С., Кушніренко І.В.,
Яблонська Т.М., Айрапетян І.А., Свердлова М.В.
Особливості перебігу соматизованих та тривожних розладів у пацієнтів з неалкогольною жировою хворобою печінки на фоні абдомінального ожиріння

| | | |
|---|-----|--|
| Ismayilova M.K. | 70 | Ісмайлова М.К. |
| Evaluation of cytokine production in patients with infertility at the stage of preparing for extracorporal fertilization | | Оцінка продукції цитокінів у пацієнтік з беспліддям на етапі підготовки до екстракорпорального запліднення |
| Kalbus O.I., Nefodov O.O. | 74 | Кальбус О.І., Нефьодов О.О. |
| The role of antibodies to acetylcholine, muscle-specific kinase, titin and SOX1 receptors in predicting severe myasthenia gravis | | Роль антитіл до рецепторів ацетилхоліну, м'язово-специфічної тирозин-кінази, титину та SOX1 у прогнозуванні тяжкого перебігу міастенії |
| Kozova I.L., Panchak O.V., Kitsera N.I., Kozovyi R.V., Dvulit Z.P., Rozhko M.M., Kovalchuk L.Ye. | 78 | Козова І.Л., Панчак О.В., Кіцера Н.І., Двуліт З.П., Рожко М.М., Ковалчук Л.Є. |
| Influence of marital status on longevity in Ukraine | | Вплив аспектів сімейного стану на довголіття в Україні |
| Korovina L.D., Zaporozhets T.M., Sulkova O.V. | 83 | Коровіна Л.Д., Запорожець Т.М., Сілкова О.В. |
| Personal characteristics and anxiety of students against the background of the COVID-19 pandemic in Ukraine | | Особистісні характеристики та тривожність студентів на фоні пандемії COVID-19 в Україні |
| Kuzminova N.V., Ivankova A.V., Lozinsky S.E., Knyazkova I.I., Ivanova E.I., Kulchytska O.M., Shkarivsky Yu.L. | 89 | Кузьмінова Н.В., Іванкова А.В., Лозинський С.Е., Князькова І.І., Іванова Е.І., Кульчицька О.М., Шкарівський Ю.Л. |
| Accordance of clinical and instrumental profile to cystatin C level in patients with stage II hypertension and frequent extrasystole | | Клініко-інструментальний профіль хворих на гіпертонічну хворобу II стадії та часту екстрасистолію відповідно до рівня цистатину С |
| Kurbanova J.F., Sultanova S.G., Hajiyeva F.R. | 94 | Курбанова Д.Ф., Султанова С.Г., Гаджиева Ф.Р. |
| Physical and psycho-emotional disorders in women with postpartum complications | | Фізичні та психоемоціональні розлади у жінок на фоні посляпологових ускладнень |
| Kuroedova V.D., Vyzhenko Ye.Ye., Stasiuk O.A., Sokolohorska-Nykina Yu.K., Petrova A.V. | 99 | Куроєдова В.Д., Виженко Є.Є., Стасюк О.А., Сокологорська-Нікіна Ю.К., Петрова А.В. |
| Cephalometric characteristics of skeletal forms of distal bite | | Цефалометрична характеристика скелетних форм дистального прикусу |
| Levytskyi A.F., Terpylovskyi Yu.R., Yaresko O.V. | 103 | Левицький А.Ф., Терпиловський Ю.Р., Ярецько О.В. |
| Biomechanical features of single-bone osteosynthesis of diaphyseal fractures in children by the titanium elastic nails | | Біомеханічні особливості single-bone остеосинтезу діафізарних переломів кісток передпліччя у дітей із застосуванням титанових еластичних стрижнів |
| Nesterenko Ye.A., Shinkaruk-Dykovytska M.M., Perlova A.V., Arshynnikov R.S., Gunas I.V. | 109 | Нестеренко Є.А., Шінкарук-Диковицька М.М., Перлова А.В., Аршинников Р.С., Гунас І.В. |
| Cephalometric parameters of basal cranial structures according to method "Cephalometrics for orthognathic surgery" in Ukrainian young men and women with orthognathic occlusion depending on the type of face | | Цефалометричні параметри базальних краніальних структур за методом «Цефалометрія для ортогнатичної хірургії» в українських юнаків і дівчат із ортогнатичним прикусом у залежності від типу обличчя |
| Pakhomova N.G., Baranets I.V., Lukianenko A.V., Leshchii N.P., Kachurovska O.B., Berezan O.I., Olefir O.I. | 113 | Пахомова Н.Г., Баранець І.В., Лук'яненко А.В., Лещій Н.П., Качуровська О.Б., Березан О.І., Олефір О.І. |
| Comprehensive rehabilitation of children with sensory and intellectual disorders | | Комплексна реабілітація дітей із сенсорними та інтелектуальними порушеннями |
| Prykhidko R.A., Dudchenko M.O., Kravtsov M.I., Zaiets S.M., Ivashchenko D.M., Chelishvili A.L., Shevchuk M.P. | 119 | Прихідко Р.А., Дудченко М.О., Кравцов М.І., Засєць С.М., Іващенко Д.М., Челішвілі А.Л., Шевчук М.П. |
| A three-stage therapeutic and diagnostic algorithm in mechanical jaundice of different genesis and the most effective mini-invasive method of its treatment | | Трьохетапний лікувально-діагностичний алгоритм при механічній жовтяниці різного генезу та найбільш ефективний мініінвазивний метод її усунення |
| Raksha-Sliusareva O.A., Tarasova I.A., Sliusarev O.A., Chernyshova O.Y., Zadorozhna V.I., Boyeva S.S., Hutarov O.I. | 125 | Ракша-Слюсарєва О.А., Тарасова І.А., Слюсарєв О.А., Чернішова О.Є., Задорожна В.І., Бояєва С.С., Гутаров О.І. |
| Cytomorphological examination in the efficiency of treatment of acute respiratory infections complicated by obstructive syndrome | | Цитоморфологічні дослідження в контролі ефективності лікування гострих респіраторних захворювань, ускладнених обструктивним синдромом |
| Rezvina K.Yu., Tkachenko P.I., Bilokon S.O., Lokhmatova N.M., Dolenko O.B., Popelo Yu.V., Korotych N.M. | 129 | Резвіна К.Ю., Ткаченко П.І., Білоконь С.О., Лохматова Н.М., Доленко О.Б., Попело Ю.В., Коротич Н.М. |
| Clinical, morphological and immunohistochemical parameters of epidermoid cysts in the neck | | Клініко-морфологічні та імунохімічні особливості епідермоїдних кіст ший |

| | | |
|--|-----|---|
| Rybachuk A.V., Tolstanov O.K., Malanchuk V.O., Mazur I.P., Fedorenko Z.P., Sumkina O.V. | 134 | Рибачук А.В., Толстанов О.К., Маланчук В.О., Мазур І.П., Федоренко З.П., Сумкіна О.В. |
| Analysis of morbidity and mortality of patients with malignant neoplasms of the lip and oral cavity in Ukraine | | Аналіз захворюваності та смертності хворих на злойкіні новоутворення губи і ротової порожнини в Україні |
| Skikevych M.G., Toropov O.A., Voloshyna L.I., Yatsenko P.I., Steblovskyi D.V., Boyko I.V., Avetikov D.S. | 141 | Скікевич М.Г., Торопов О.А., Волошина Л.І., Яценко П.І., Стебловський Д.В., Бойко І.В., Аветіков Д.С. |
| Analysis of clinical parameters of scar tissue of the scalp and neck depending on the circadian rhythm of the patient | | Аналіз клінічних параметрів рубцевозміненої тканини голови та шиї в залежності від циркадного ритму пацієнта |
| Skrupnyk I.M., Maslova G.S., Lymanets T.V. | 146 | Скрипник І.М., Маслова Г.С., Лиманець Т.В. |
| Overweight and obesity as risk factors for chemotherapy-induced hepatotoxicity in patients with acute lymphoblastic leukemia | | Надлишкова вага та ожиріння як фактор ризику цитостатик-індукованих уражень печінки у хворих на гостру лімфобластну лейкемію |
| Sluchevska O.O., Pavlenko O.V., Mochalov I.O., Kryvtsova M.V., Tsaryk V.V., Karbovanets O.I. | 151 | Случевська О.О., Павленко О.В., Мочалов Ю.О., Кривцова М.В., Царик В.В., Карбованець О.І. |
| Peculiarities of the periodontal pockets microbiota during acute duration of generalized periodontitis | | Особливості мікробіоти пародонтальних кишень при гостром перебігу генералізованого пародонтиту |
| Stepanenko R.L., Dudchenko M.O., Stepanenko V.I., Vasyljeva K.V., Kameniev V.I., Svirid S.G., Litinska T.O. | 156 | Степаненко Р.Л., Дудченко М.О., Степаненко В.І., Васильєва К.В., Каменєв В.І., Свірид С.Г., Літінська Т.О. |
| Lipid parameters before and after immunobiological therapy of patients with psoriasis | | Показники ліпідів до та після проведення імунобіологічної терапії у хворих на псоріаз |
| Stoyanov O.M., Vastyanov R.S., Myronov O.O., Kalashnikov V.I., Babienko V.V., Hruzevskiy O.A., Turchin M.I. | 162 | Стоянов О.М., Вастьянов Р.С., Миронов О.О., Калащенков В.Й., Бабінко В.В., Грузевський О.А., Турчин М.І. |
| Vegetative system pathogenetic role in chronic brain ischemia, cerebral hemodynamics disorders and autonomous dysregulation | | Патогенетична роль вегетативної системи при формуванні хронічної ішемії мозку, порушень церебральної гемодинаміки та автономної регуляції |
| Tyazhelov A.A., Karpinska O.D., Karpinsky M.Yu., Klymovytskyi V.H., Goncharova L.D., Yurchenko D.O. | 168 | Тяжелов О.А., Карпінська О.Д., Карпінський М.Ю., Климовицький В.Г., Гончарова Л.Д., Юрченко Д.О. |
| X-ray characteristics of the dysplastic hip joint in adults before and after endoprosthetic surgery | | Рентгенологічна характеристика диспластичного стегнового суглобу у дорослих до та після операції ендопротезування |
| Khasawneh A.R., Dmytrenko S.V., Shevchuk Yu.G., Kondratuk A.I., Kryvonis T.G. | 174 | Хасавнек А.Р., Дмитренко С.В., Шевчук Ю.Г., Кондратюк А.І., Кривоніс Т.Г. |
| Discriminant models of seborheic dermatitis possibilities of occurrence and course in men and women depending on the characteristics of anthroposomatotypological parameters | | Дискримінантні моделі можливості виникнення та особливостей перебігу себорейного дерматиту у чоловіків та жінок у залежності від особливостей антро-соматотипологічних показників |
| Shaposhnyk O.A., Prykhodko N.P., Savchenko L.V., Shevchenko T.I., Sorokina S.I., Yakymyshyna L.I., Kudria I.P. | 178 | Шапошник О.А., Приходько Н.П., Савченко Л.В., Шевченко Т.І., Сорокіна С.І., Якимишина Л.І., Кудря І.П. |
| Clinical and diagnostic aspects of managing patients with valvular heart disease | | Клініко-діагностичні аспекти ведення пацієнтів із захворюваннями клапанів серця |
| Shevchuk I.M., Novitsky O.V., Shapoval A.L., Sadovy I.Ya., Sverstyuk A.S., Snizhko S.S. | 183 | Шевчук І.М., Новицький О.В., Шаповал А.Л., Садовий І.Я., Сверстюк А.С., Сніжко С.С. |
| Using ultrasound diagnostics and magnetic resonance tomography in surgical treatment of complicated forms of acute paraproctitis | | Застосування ультразвукової діагностики і магнітно-резонансної томографії у хірургічному лікуванні складних форм гострого парапроктиту |

EXPERIMENTAL MEDICINE

| | |
|--|-----|
| Appelhans O.L., Kozishkurt O.V., Gritsenko K.S., Boruta N.V., Shabliy T.P., Hruzevskiy O.A., Naidich O.V. | 189 |
| Molecular and genetic characteristics of adenovirus as causes of acute diarrhea in the people of southern Ukraine | |
| Dienha O.V., Osadcha A.O., Makarenko O.A., Babenia H.O., Horokhivsky V.N., Shnaider S.A. | 194 |
| Experimental evaluation of the effectiveness of the use of a drugs complex developed for the prevention of major dental diseases in young athletes | |

ЕКСПЕРИМЕНТАЛЬНА МЕДИЦИНА

| | |
|---|-----|
| Appelman O.L., Kozishkurt O.V., Gritsenko K.S., Boruta N.V., Shabliy T.P., Gruzhevskyi O.A., Naidich O.V. | 189 |
| Молекулярно-генетичні характеристики адено-вірусів, як збудників гострих діарей у жителів півдня України | |
| Деньга О.В., Осадча А.О., Макаренко О.А., Бабеня Г.О., Горохівський В.Н., Шнейдер С.А. | 194 |
| Експериментальна оцінка ефективності використання комплексу засобів, розробленого для профілактики основних стоматологічних захворювань у дітей-спортсменів | |

- Yeroshenko G.A., Grygorenko A.S., Shevchenko K.V., Lysachenko O.D., Riabushko O.B., Pyvovar N.M., Klepets O.V.**
Influence of food additives complex on the morphology of villi of the rats' duodenum mucosa
- Kvitnitskaya-Ryzhova T.Yu., Lugovskyi S.P., Klymenko P.P., Mykhalskiy S.A., Malyshova S.P., Tkachenko O.S., Kosiakova H.V.**
The influence of age factor on changes in morphometric indices of rat pancreas in modeling of insulin resistance and its correction with N-stearoylethanolamine
- Kravets N.Ya., Klumnyk S.I., Romanyuk L.B., Borak V.P.**
Biofilm-forming properties of pathogenic microorganisms in children with recurrent tonsillitis
- Mykytenko A.O., Akimov O.Ye., Yeroshenko G.A., Neporada K.S.**
Influence of transcription factor κB on remodeling of extracellular matrix of rat liver under conditions of chronic alcohol intoxication
- Nosivets D.S., Shevchenko T.M., Govorukha O.Yu.**
Influence of non-steroidal anti-inflammatory drugs on movement-research behavior of animals under experimental secondary osteoarthritis
- Pivtorak V.I., Sydorenko B.V., Monastyrskyi V.M., Pivtorak, K.V. Bulko M.P.**
Efficacy of an experimental model of non-alcoholic fatty liver disease based on a high-fat diet with cholesterol
- Safonov R.A., Prokopiuk V.Yu., Prokopiuk O.V., Lazurenko V.V., Tishchenko O.M., Ovcharenko O.B.**
Comparison of the effect of different mesh implants on the reproductive system in experimental models of gynecological surgeries
- Stepanchuk A.P., Fedorchenco I.L., Pryshliak A.M., Piliuhin A.V., Shepitko K.V.**
The normal architectonics of the greater omentum vasculature in elderly people of both gender
- Tiron O.I., Stetsenko A.V., Yatsyna O.I., Zayats L.M., Kolotvin A.O., Shumilina K.S.**
The morphological changes of the white rats' thyroid gland 21 days after experimental thermal burn injury under NaCl systemic administration
- Frenkel Yu.D., Chernov V.S., Kharchenko O.V., Larycheva O.M., Chebotar L.D., Pshychenko V.V., Kostenko V.O.**
Main pathomorphological changes of liver tissues in experimental metabolic syndrome
- Khmara T.V., Stelmakh G.Ya., Garvasiuk O.V., Biriuk I.G., Drachuk V.M.**
Peculiarities of innervation of muscles and skin of the anterior and lateral abdominal walls in human fetuses
- Yachmin A.I., Yeroshenko G.A., Shevchenko K.V., Hapon S.V., Vatsenko A.V., Ulanovska-Tsyba N.A., Sokolenko V.M.**
Ultrastructural characteristics of the rat gastric fundic wall after the impact of the complex of food additives
- 199 Єрошенко Г.А., Григоренко А.С., Шевченко К.В., Лисаченко О.Д., Рябушко О.Б., Пивовар Н.М., Клепець О.В.**
Вплив комплексу харчових добавок на морфологію ворсин слизової оболонки 12-палої кишки щурів
- 204 Квітницька-Рижова Т.Ю., Луговський С.П., Клименко П.П., Михальський С.А., Малишева С.П., Ткаченко О.С., Косякова Н.В.**
Вплив фактора віку на зміни морфометричних показників підшлункової залози шура при моделювання інсулінерезистентності та її корекції N-стеароїлєтаноламіном
- 210 Кравець Н.Я., Климнюк С.І., Романюк Л.Б., Борак В.П.**
Плівкоутворюючі властивості умовно-патогенних мікроорганізмів у дітей, хворих на рекурентний тонзіліт
- 214 Микитенко А.О., Акімов О.Є., Єрошенко Г.А., Непорада К.С.**
Вплив транскрипційного фактору κB на ремоделювання екстрацелюлярного матриксу печінки щурів за умов хронічної алкогольної інтоксикації
- 218 Носівець Д.С., Шевченко Т.М., Говоруха О.Ю.**
Вплив ненаркотичних анальгетиків на рухово-дослідницьку поведінку тварин за умов експериментального вторинного остеоартрозу
- 222 Півторак В.І., Сидоренко Б.В., Монастирський В.М., Півторак К.В., Булько М.П.**
Дієвість експериментальної моделі неалкогольної жирової хвороби печінки на основі високожирової дієти та холестерину
- 227 Сафонов Р.А., Прокопюк В.Ю., Прокопюк О.В., Лазуренко В.В., Тіщенко О.М., Овчаренко О.Б.**
Порівняння впливу різних сітчастих імплантів на статеву систему в експериментальних моделях гінекологічних операцій
- 232 Степанчук А.П., Федорченко І.Л., Пришляк А.М., Піллогін А.В., Шепітько К.В.**
Архітектоніка судин великого чепця в жінок і чоловіків похилого віку в нормі
- 237 Тірон О.І., Стеценко А.В., Яцина О.І., Заяць Л.М., Колотвін А.О., Шуміліна К.С.**
Морфологічні зміни щитоподібної залози білих щурів через 21 добу після експериментального термічного опіку при системному введенні NaCl
- 242 Френкель Ю.Д., Черно В.С., Харченко О.В., Ларичева О.М., Чеботар Л.Д., Пшиченко В.В., Костенко В.О.**
Основні патоморфологічні зміни тканин печінки при експериментальному метаболічному синдромі
- 247 Хмара Т.В., Стельмах Г.Я., Гарвасюк О.В., Бірюк І.Г., Драчук В.М.**
Особливості іннервації м'язів і шкіри передньої і бічних стінок живота у плодів людини
- 252 Ячмінь А.І., Єрошенко Г.А., Шевченко К.В., Гапон С.В., Ваценко А.В., Улановська-Циба Н.А., Соколенко В.М.**
Ультраструктурна характеристика фундального відділу стінки шлунку щурів після дії комплексу харчових добавок

EVENTS AND DATES

To the blessed memory of professor Chaikovskyi Yurii Bohdanovich

НОВІ ДАТИ

256 Пам'яті професора Чайковського Юрія Богдановича

A.P. Stepanchuk, I.L. Fedorchenko, A.M. Pryshliak¹, A.V. Piliuhin, K.V. Shepitko²

Poltava State Medical University, Poltava

¹I. Horbachevsky Ternopil National Medical University, Ternopil

²Municipal non-profit enterprise "Kotelya Hospital of planned treatment" Kotelya

THE NORMAL ARCHITECTONICS OF THE GREATER OMENTUM VASCULATURE IN ELDERLY PEOPLE OF BOTH GENDER

e-mail: fedorchenkoigor@ukr.net

Variability of topography of arteries and veins of the greater omentum, taken from 20 cadavers of elderly people was investigated by the methods of anatomical preparation and morphometry. The right and left gastro-omental arteries form the superior arterial arch in 15 of 18 cases, and did not anastomose in 3 cases. The inferior arterial arch was present in 1 of 20 cases. The left gastro-omental artery was absent in 2 of 20 cases. The central artery has a greater length and outer diameter both in women and men. The veins, in comparison with the arteries, have a larger outer diameter and one vein accompanies the artery of the same name. The zone of the central, and in its absence – the right and left medial arteries of the greater omentum can be used for obtaining the graft for transplantation in the quadrangular and triangular shapes of the greater omentum of the elderly people.

Key words: human greater omentum vasculature, morphometry, topography of arteries, gastro-omental arteries, right gastro-omental arteries, left gastro-omental arteries

А.П. Степанчук, І.Л. Федорченко, А.М. Пришляк, А.В. Пілюгін, К.В. Шепітько

АРХІТЕКТОНІКА СУДИН ВЕЛИКОГО ЧЕПЦЯ В ЖІНОК І ЧОЛОВІКІВ ПОХИЛОГО ВІКУ В НОРМІ

Методами анатомічного препарування та морфометрії досліджена варіабельність топографії артерій і вен великих чепців від 20 (по 10 чоловічої і жіночої статі) трупів людей похилого віку. Права і ліва шлунково-чепцеві артерії утворюють верхню артеріальну дугу в 15 із 18 випадків, а в 3 випадках артерії не з'єднувалися. Нижня артеріальна дуга була наявна в 1 із 20 випадків. Ліва шлунково-чепцева артерія в 2 із 20 випадків була відсутня. Центральна артерія має більшу довжину і зовнішній діаметр у жінок і чоловіків. Вени, в порівнянні з артеріями, мають більший зовнішній діаметр і в кількості однієї вени супроводжують одноіменну артерію. Венозна кров від великого чепця відтікає в систему ворітної вени через верхню брижову і селезінкову вени. Зона центральної, а при її відсутності правої і лівої медіальних артерій великого чепця може використовуватися для трансплантації при чотирикутній і трикутній формах великого чепця в людей похилого віку.

Ключові слова: судини великого чепця людини, морфометрія, топографія артерій, шлунково-чепцеві артерії, права шлунково-чепцева артерія, ліва шлунково-чепцева артерія.

The study is a part of the research project "Morphofunctional study of the internal organs of humans and laboratory animals in different aspects of the experimental medicine"; state registration No. 0121U108258

Rare cases of primary torsion of the omentum and idiopathic segmental omental infarction are treated by surgical laparoscopic removal (omentectomy) within healthy tissues to avoid complications, gastric vein thrombosis and portal vein thrombosis of the liver [2, 3, 7, 8]. To optimize the surgical procedure of excision of the autograft from the large omentum, the surgeon must be guided by the variability of its vascular architecture and vascular bifurcation standards. Current domestic publications do not elucidate data on the topography, length and diameters of vessels of the human greater omentum. At the same time, since no analysis of data on their normal number and branches in the omentum has been found, our study was aimed at this issue. In the literature, there are differences and few data on the individual and age characteristics of the vessels of the human greater omentum [9, 10]. In case of unsuccessful conservative treatment of primary or secondary inflammation of the greater omentum, its resection is indicated. Being involved in the intercurrent process, the greater omentum can twist, and with a change in intra-abdominal pressure and hyperperistalsis of the small intestine, the right part of the greater omentum (with its multi-lobed form) can wrap, which contributes to disruption of its blood supply due to vascular compression [8]. All of the above often puts the surgeon in front of the need for surgery on a large omentum, which requires knowledge of variations in topography and the number of vessels in the omentum in humans.

The purpose of the study was to establish the topographic and morphometric variability of the arteries and veins of the greater omentum from elderly women and men.

Materials and methods. The arteries and veins of the greater omentum from 10 male cadavers and 10 female cadavers have been studied. The deceased were elderly people without intra-abdominal lesions. Vasculature of the greater omentum was examined during the autopsy in accordance with the Cooperation Agreement with the Pathology Department of the CE "O.F. Maltsev Poltava Regional Clinical Hospital PRC" and Poltava Regional Autopsy Bureau. The method of anatomical preparation of vessels on

the greater omentum with moderately and excessively developed adipose tissue has been used. Subsequently, to examine the vessels, the preparations of the greater omenta were placed on the negatoscope, then photographed and the value of the parameters of the length and outer diameter of vessels and branches at the point of their origination from the main vessels was measured using a caliper. The study was performed in compliance with the basic bioethical provisions of the Council of Europe Convention on Human Rights and Biomedicine (04.04.1997), the Helsinki Declaration of the World Medical Association on the ethical principles of human research (1964–2008), and the Order of the Ministry of Health of Ukraine No. 690 as of September 23, 2009.

To determine the arithmetic mean and mean deviation of the parameters of the length and outer diameter of the vessels and their branches, statistical processing of the obtained data was performed using the Microsoft Excel 2010. The nonparametric Mann-Whitney test was used to determine the statistical significance of differences between the groups. The difference was considered statistically significant at $p < 0.05$.

Results of the study and their discussion. Macroscopically, the right gastro-omental artery was detected in all 20 preparations and the left gastro-omental artery was detected in 90 % (18 cases out of 20) on the secured part of the greater omentum. The right and left gastro-omental arteries anastomose with each other on the secured part of the greater omentum in the area of the greater curvature of the stomach at the border of the middle and upper thirds of its curvature in 83 % (15 cases out of 18), forming the superior arterial arch of the greater omentum. No anastomosis with the left and right gastro-omental arteries was noted in 3 (16 %) out of 18 cases in the presence of the above part of the greater omentum. Numerous branches originated from both arteries and run to the greater curvature of the stomach. On the greater omentum from female and male cadavers, the length of the right gastro-omental artery ranged from 14.5 to 38 cm and from 10.6 to 26 cm, respectively, and its diameter was 0.16 - 0.38 cm and 0.2–0.32 cm, respectively. In addition, 3 to 12 branches commonly arise from the right gastro-omental artery. These branches ascended by 2–4 cm along the anterior wall of the stomach, entering its depth. Subsequently, we have found that both female and male greater omentum was supplied with blood by the omental arteries, bifurcated from the right or left gastric arteries, when they were located in the area of the lesser curvature of the stomach in 25 % (5 cases out of 20). The analysis of our own studies enabled to distinguish the following omental branches of the right gastro-omental artery: the central artery, the right medial artery, the right intermediate artery, the right lateral artery, the left extra artery. The central omental artery was found in 85 % (17 cases out of 20) and was located in the middle of the greater omentum (fig. 1).



Fig. 1. Anatomical preparation of the greater omentum of elderly people. 1 – the right gastro-omental artery; 2 – the stomach; 3 – the left gastro-omental artery; 4 – the left medial artery; 5 – the central artery; 6 – the right medial artery; 7 – the right lateral artery.

respectively, and the diameter ranged from 0.11 to 0.15 cm. The right medial artery was located by 2–3 cm to the right of the central omental artery, arose at an angle of 60°–80° from the right gastro-omental artery, running to the lower edge of the unsecured part of the omentum. The right medial artery gave rise to the branches in the middle third of the length of the omentum, and its terminal branches arose from the lower edge of the unsecured part of the greater omentum. In 2 cases, the right medial artery anastomosed with the right lateral artery in the right area of the unsecured part of the greater omentum in its middle third.

The right intermediate artery was found in 45 % (9 cases out of 20) on the greater omentum from the elderly subjects. On the female and male greater omenta, the length of the right intermediate artery

This artery originated from the right gastro-omental artery at an angle of 85°–90°, running to the lower edge of the unsecured part of the omentum. It was observed that the right lateral branch of the central omental artery anastomosed with the right medial omental artery (1 case), and in two cases the left lateral branch anastomosed with the left lateral omental artery (the branch of the left gastro-omental artery) and with the left medial omental artery (the branch of the left gastro-omental artery) in one case. On the greater omentum from female and male cadavers, the length of the central omental artery ranged from 16.2 to 30 cm and from 17 to 25.7 cm, respectively, and the diameter ranged from 0.12 to 0.22 cm and 0.13 to 0.29 cm, respectively.

On the greater omentum from female and male cadavers, the length of the right medial artery ranged from 6 to 31.5 cm, and from 17 to 25.7 cm, respectively. The right medial artery was located by 2–3 cm to the right of the central omental artery, arose at an angle of 60°–80° from the right gastro-omental artery, running to the lower edge of the unsecured part of the omentum. The right medial artery gave rise to the branches in the middle third of the length of the omentum, and its terminal branches arose from the lower edge of the unsecured part of the greater omentum. In 2 cases, the right medial artery anastomosed with the right lateral artery in the right area of the unsecured part of the greater omentum in its middle third.

ranged from 13 to 27.3 cm and 13.2 to 22 cm, respectively, and the diameter was 0.11 cm and 0.1 to 0.14 cm, respectively.

The right lateral artery was found in 70 % (14 cases out of 20) on the greater omentum from the elderly subjects. On the female and male greater omenta, the length of the right lateral artery ranged from 7.5 to 18.8 cm and 8 to 16.5 cm, respectively, and the diameter was 0.1–0.11 cm and 0, 1–0.12 cm, respectively.

In two cases, on the female greater omentum it was observed that the right gastro-omental artery gave rise to the left extra arteries. In the first case, the left extra artery was located on the left half of the greater omentum by 3 cm laterally to the central artery and anastomosed with the left lateral artery (the branch of the left gastroesophageal artery). In the second case, the central omental artery was absent, and the left extra artery ran along the left half of the entire length of the omentum to its lower edge, where it ended. The length of the left extra arteries ranged from 18 to 21.5 cm, and the diameter was 0.11 to 0.13 cm.

The left gastro-omental artery was absent on the anterior plate of the greater omentum in 10 % (per 1 case in women and men). Being located in the depth of the posterior surface of the secured part of the great omentum, the left gastro-omental artery also gave rise to 7–13 gastric branches. On the greater omenta of female and male cadavers, the length of the left gastro-omental artery ranged from 11 to 20.5 cm and 8 to 16 cm, respectively, and the diameter ranged from 0.12 to 0.21 cm and 0.13 to 0.24 cm, respectively. Our own observations enabled to distinguish the following branches of the left gastro-omental artery: left medial artery, left intermediate artery, left lateral artery. The left medial artery was found in 89 % (16 cases out of 18), the left intermediate artery in 33 % (6 cases out of 18), and the left lateral artery in 89 % (16 cases out of 18). These branches arose at an angle of 45°–70° from the left gastro-omental artery, branching into numerous fine arterial vessels in the left upper and middle parts of the greater omentum. The left medial, intermediate and lateral arteries do not reach the lower edge of the omentum.

On the greater omenta from female and male cadavers, the length of the left medial omental artery ranged from 8.5 to 29.5 cm and 8.5 to 28.2 cm, respectively, and the diameter ranged from 0.12 to 0.18 cm and 0.11 to 0.14 cm, respectively.

Mostly, in 77 % (14 cases out of 18), the left intermediate artery were absent on the greater omenta. If present on the female and male greater omenta, the arteries length ranged from 14.7 to 21 cm and 8.5 to 16.5 cm, respectively, and the diameter ranged from 0.11 to 0.12 cm and 0.11 to 0.12 cm, respectively.

On the female and male greater omenta, the length of the left lateral artery ranged from 6 to 25 cm and 7 to 27.2 cm, respectively, and its diameter ranged from 0.1 cm to 0.2 cm and 0.1 to 0.22 cm, respectively. In one of 18 cases, the left lateral artery anastomosed with the left medial artery in the left upper third of the unsecured part of the greater omentum. Notably, in two cases, when the left gastro-omental artery was located on the posterior surface of the secured part of the greater omentum, it gave rise, except for the left lateral artery, to no other branches.

In the first case, when originated from the left gastro-omental artery, the left lateral artery was located by 2 cm inward along the left edge of the anterior plate of the unsecured part of the greater omentum and branched into numerous fine arterial branches. Subsequently, the left extra artery (the branch of the right gastro-omental artery) entered the artery at the level of the left middle part of the greater omentum,

and then below at a distance of 3 cm the left lateral artery turned to the right and formed anastomosis with the central artery (the branch of the right gastro-omental artery) in the lower third of the unsecured part of the omentum.

In the second case, the left lateral artery was located medially and vertically of the left part of the anterior plate of the greater omentum, where it gave rise to two branches, ending in the depth of the left edge of the omentum.

Then the artery reached the middle of the lower part of the omentum and anastomosed with the right lateral artery (the branch of the right gastro-omental artery), forming the inferior arterial arch of the omentum on its anterior plate. Notably, no central artery was found on the greater omentum (fig. 2) and 3 omental branches originated from the left gastric artery were detected.



Fig. 2. The arteries of the elderly human greater omentum. 1 – the right gastro-omental artery; 2 – the stomach; 3 – omental branches of the left gastric artery; 4 – the left lateral artery; 5 – inferior arterial arch of the greater omentum; 6 – the right lateral artery.

These branches descended to the left side of the anterior wall of the stomach, then anastomosed with each other in the left upper third of the secured part of the greater omentum, giving rise to the arterial branch, which bifurcated in the left middle part of the anterior plate of the greater omentum and supplied blood to it.

In addition to the above-mentioned branches, in 2 out of 20 cases the upper left part of the greater omentum was supplied with blood by the omental-splenic artery, originated from the splenic artery.

The resulting morphometric data showed that the central artery of the greater omentum from the elderly women and men had the highest value of the outer diameter and the length compared to the arterial branches listed above (table 1).

Table 1

The parameters of the vasculature of the human greater omentum M±m

| Vessels | Parameters | | | |
|-----------------------------|--------------|----------------|--------------|----------------|
| | Diameter, cm | | Length, cm | |
| | women | men | women | men |
| Right gastro-omental artery | 0.232±0.071 | 0.25±0.071* | 21.16±13.65 | 16.35±8.301* |
| Central artery | 0.138±0.029 | 0.177±0.095* | 21.48±8.131 | 22.45±3.511* |
| Right medial artery | 0.12±0.013 | 0.12±0.016* | 18.43±13.08 | 19.3±0.019* |
| Right intermediate artery | 0.11±0 | 0.12±0.016* | 20±11.03 | 18.54±6.118* |
| Right lateral artery | 0.102±0.0027 | 0.108±0.011* | 12.96±8.126 | 12.978±5.72* |
| Left gastro-omental artery | 0.176±0.054 | 0.193±0.048* | 13.3±5.481 | 11.91±4.065* |
| Left medial artery | 0.134±0.017 | 0.124±0.013* | 16.729±15.03 | 17.87±9.21* |
| Left intermediate artery | 0.115±0.003 | 0.115±0.008* | 17.85±2.1 | 13.85±5.14* |
| Left lateral artery | 0.116±0.0517 | 0.125±0.0563* | 14.322±10.62 | 14.457±10.907* |
| Right gastro-omental vein | 0.267±0.105 | 0.28±0.075* | 21.16±13.65 | 16.35±8.30* |
| Central vein | 0.1533±0.021 | 0.1938±0.106* | 21.48±8.131 | 22.45±3.511* |
| Right medial vein | 0.136±0.014 | 0.144±0.019* | 18.43±13.08 | 19.36±0.019* |
| Right intermediate vein | 0.127±0.003 | 0.136±0.009* | 20±0.003 | 18.54±6.118* |
| Right lateral vein | 0.124±0.0104 | 0.1267±0.019* | 12.96±8.126 | 12.978±5.72* |
| Left gastro-omental vein | 0.215±0.07 | 0.218±0.046* | 13.3±5.481 | 11.91±4.065* |
| Left medial vein | 0.151±0.019 | 0.1411±0.017* | 16.729±15.03 | 17.87±9.213* |
| Left intermediate vein | 0.135±0.003 | 0.1325±0.007* | 17.85±2.1 | 13.85±5.14* |
| Left lateral vein | 0.1344±0.059 | 0.1486±0.0645* | 14.322±10.62 | 14.457±10.907* |

*— the reliable difference between women and men makes up $p<0.05$.

In our opinion, such anatomical features of the vascular topography, variations of branching and formation of anastomosis are of practical importance during surgical interventions on the greater omenta in their pathology and in reconstructive surgery, when a graft is cut from the greater omentum for transplantation to an organ with defects.

There are facts in the literature that indicate that the greater omentum of the elderly people has a quadrangular, triangular and multi-lobe shape. The quadrangular omentum had a width of 25 to 35 cm, which exceeded their length (17.5–27 cm). Triangular-shaped greater omentum are rare and have a wide base and a narrow tapered lower edge, and their length is from 25 to 37 cm, width is 22.5–29.4 cm. The length of irregularly shaped omentum ranged from 14 to 30 cm, and a width of 28 cm (the smallest) to 52 cm (the largest). The quadrangular shape of the greater omentum is more favorable for obtaining a graft with elongation for transplantation to organs with a large defect [6]. We also verified the above shapes of the greater omentum of the elderly people in 45 % (9 cases out of 20) as quadrangular, in 15 % (3 cases out of 20) as triangular and in 40 % (8 cases out of 20) of irregular shape with two or more lobes.

The analysis of our own studies showed that the left and right halves of the quadrangular greater omentum had the same blood supply, which was provided by the right and left gastro-omental arteries and their branches. In the greater omental of irregular shape with two lobes more favorable blood supply was found in the right half of the greater omentum. The upper left part of the greater omentum had a more optimal blood supply in the presence of the omental-splenic artery. From the triangular-shaped greater omentum it is possible to obtain a short autograft and a longer one from the quadrangular-shaped greater omentum.

In our opinion, to obtain the graft from the greater omentum, both its shape and the length and diameter of the vessels should be taken into account. The central artery (a branch of the right gastro-omental artery) of the greater omentum from female and male cadavers had a larger length and outer diameter compared to the studied arteries and in its absence the right and left medial arteries in men and the left and right intermediate arteries in women.

The right gastro-omental artery is a branch of the gastroduodenal artery, and the left gastro-omental artery arises from the splenic artery [1, 4]. The facts of anastomoses described by us are confirmed by the data of other authors [4, 5]. Stembre N. [9] reports that the average length of the right gastro-omental artery is 244.3 ± 34.4 mm, and the diameter is 2.68 ± 0.39 mm. The diameter of the right gastro-omental artery is ≥ 3 mm at the beginning and 1.5–2 mm at the level of the middle of the greater curvature of the stomach [10].

Some authors [6] highlight the omental branches of the posterior plate of the unsecured part of the greater omentum anastomose with the middle colic and inferior pancreatic arteries. However, our studies found no anastomoses. Instead, we observed anastomoses between the arterial branches of the anterior and posterior plates when conjoined.

The pathway of the veins of the greater omentum was similar to the pathway of the arteries. Each artery was accompanied by one vein. Veins were formed from the conjoining of numerous fine branches that arose from the depth of the lower and middle areas of the unsecured part of the greater omentum. In these areas, the venous branches had a transverse pathway, they conjoined and gave rise to the central, right medial, intermediate and lateral veins of the right half of the omentum. Respectively left medial, intermediate and lateral veins were formed in the left part of the greater omentum. The outer diameter of the veins is wider than the diameter of the arteries. Table 1 presents the average values of the parameters of the veins. Then the veins ascended along with the anterior plate of the unsecured part of the greater omentum to the upper edge of the secured part of the omentum and formed the right and left gastro-omental veins, which were located parallel to the greater curvature of the stomach. The right gastro-omental vein flowed into the superior mesenteric vein, and the left one into the splenic vein.

Conclusions

1. In the elderly people, the blood supply to the greater omentum is provided by the right and left gastro-omental arteries. These arteries anastomose and form the superior arterial arch on the secured part of the greater omentum in 15 out of 18 cases and in 3 out of 18 cases the arteries did not anastomose at all. The left gastro-omental artery in 2 of 20 cases was absent on the anterior plate of the greater omentum. The lower edge of the unsecured part of the omentum had the arterial arch in 1 out of 20 cases.

2. The right gastro-omental artery give rise to the central, right medial, intermediate and lateral arteries and left extra artery (in 2 out of 20 cases). The central artery has a length of 21.49 ± 8.1 cm, an outer diameter of 0.14 ± 0.029 cm in women and 22.45 ± 3.5 cm and 0.17 ± 0.09 cm in men, respectively. The branches of the right gastro-omental artery supply blood to the right half of the greater omentum and reach the lower edge of the unsecured part of the omentum. The left extra artery branches supply blood to the left half of the omentum. The left gastro-omental artery bifurcates into the left medial, intermediate and lateral arteries. These arteries branch into the depth of the left half of the omentum and supply blood to it without reaching the lower edge of the unsecured part of the omentum. The upper left area of the omentum can be supplied with blood by the splenic-omental artery, which arises from the splenic artery in 2 out of 20 cases.

3. Veins, compared to arteries, have bigger external diameter and one vein accompanies an artery of the same name. The topography of the omental veins is similar the topography of the omental arteries. Venous blood outflows from the greater omentum into the portal vein system through the superior mesenteric and splenic veins.

4. The greater omenta of quadrangular, triangular and irregular shape with two or more lobes were found in 45 %, 15 % and 40 %, respectively. The area of the central artery, and in case of its absence – the right and left medial arteries of the greater omentum is more favorable for obtaining the graft for transplantation in the quadrangular and triangular shapes of the greater omentum of the elderly people.

References

1. Guseynov TS, Guseynova ST, Suleymanova RG, Atayeva DA, Khalilov MA. Osobennosti anatomii bolshogo salnika. 2018; [tsitovano 2021 Marta 11]; (6): 16–8. doi:10.24411/2075-4094-2018-16276 [in Russian]
2. Kamynin YuV, Proshchenko YaN, Fayzulayev VKh, Zuyev ME. Idiopaticheskiy segmentarnyy infarkt bolshogo salnika u rebenka. Detskaya khirurgiya. 2015; 19 (4): 51–3. [in Russian]
3. Konoplitskiy VS, Pogoriliy VV, Fomin OO, Ksonz IV, Yakimenko OG, Gorbatuk LV, ta in. Torsiyna patologiya velikogo cheptsa u ditey: oglyad literaturi ta vlasni klinichni sposterezheniya. Khirurgiya dityachogo viku. 2019; 2 (63): 84–91. DOI 10.15574 / PS.2019.63.84 [in Ukrainian]
4. Shevlyuk NN, Khalikova LV, Khalikov AA. Morfofunktionalnaya kharakteristika bolshogo salnika. Zhurnal anatomii i histologii. 2020; 9 (2): 90–9. [in Russian]

5. Fedorchenko IL, Stepanchuk AP. Zovnishnya budova velikogo cheptsa lyudini u vikovomu aspekti. Aktualni problemi suchasnoyi meditsini: Visnik Ukrayinskoj medichnoyi stomatologichnoyi akademiyi. 2020; 20 (3 (71)): 222–8. [in Ukrainian]
6. Khodov NA. Klinicheskaya anatomiya bolshogo salnika. Molodoy uchonyy. 2018 Dekabr; (51): 76–80. [in Russian]
7. Cupra R, Farhat W, Ammar H, Azzaza M, Lagha S, Cheikh Y, et al. Idiopathic segmental infarction of the omentum mimicking acute appendicitis: A case report. Int J Surg Case Rep. 2019; 22 (5):18–20.
8. Kataoka J, Nitta T, Ota M, Takashima Y, Yokota Y, Fujii K, et al. Laparoscopic omentectomy in primary torsion of the omentum: report of a case. Surg Case Rep. 2019; 5 (1): 76–8. <https://doi.org/10.1186/s40792-019-0618-5>
9. Settembre N, Labrousse M, Magnan PE, Branchereau A, Champsaur P, Bussani R, et al. Surgical anatomy of the right gastro-omenta artery: a study on 100 cadaver dissections. Surg Radiol Anat. 2018 Apr; 40 (4): 415–422. <https://doi.org/10.1007/s00276-017-1951-7>
10. Suma H. Gastroepiploic artery graft in coronary artery bypass grafting. Ann Cardiothorac Surg. 2013 Jul; 2 (4): 493–8.

Стаття надійшла 22.05.2021 р.

DOI 10.26724/2079-8334-2022-2-80-237-241

UDC 616.441:599.323.4:615.459

О.І. Тірон, А.В. Стєценко, О.І. Яцина¹, Л.М. Заяць², А.О. Колотвін, К.С. Шуміліна

Odessa National Medical University, Odessa;¹ National Cancer Institute, Kyiv

²Ivano-Frankivsk National Medical University, Ivano-Frankivsk

THE MORPHOLOGICAL CHANGES OF THE WHITE RATS' THYROID GLAND 21 DAYS AFTER EXPERIMENTAL THERMAL BURN INJURY UNDER NaCl SYSTEMIC ADMINISTRATION

e-mail: chekina.o@ukr.net

The histological features of the experimental animals' thyroid gland 21 days after modelled burn injury under NaCl systemic administration can be described as adaptive and compensatory processes in the follicular wall, vascular and stromal components. Most of the thyroid follicles were round or oval in shape, some of them were overstretched and filled with dense colloid. The follicular epithelial cells were flattened, contained pyknotic nuclei surrounded by thin layer of cytoplasm. In some regions it was observed the desquamation of the thyrocytes within the follicular lumen. The electron microscopic examination of the gland 21 days after modelled burn confirmed the changes that had been previously established by the analysis of histological specimens. The flattened thyrocytes contained osmophilic nuclei with heterochromatin. Their cytoplasm was electron lucent and contained organelles exhibiting the signs of destruction. The microvilli were observed on the apical domain of thyrocytes. Along with the follicles with altered histological structure (which indicates an imbalance in the synthesis and excretion of hormones) there were observed the follicles with intact wall and unaffected blood supply. The ultrastructure of the latter indicates active protein-synthetizing and protein-secreting processes.

Keywords: burn injury, thyroid gland, light microscopy, electron microscopy.

О.І. Тірон, А.В. Стєценко, О.І. Яцина, Л.М. Заяць, А.О. Колотвін, К.С. Шуміліна МОРФОЛОГІЧНІ ЗМІНИ ЩИТОПОДІБНОЇ ЗАЛОЗИ БІЛИХ ЩУРІВ ЧЕРЕЗ 21 ДОБУ ПІСЛЯ ЕКСПЕРИМЕНТАЛЬНОГО ТЕРМІЧНОГО ОПІКУ ПРИ СИСТЕМНОМУ ВВЕДЕННІ NACL

Гістологічні особливості щитовидної залози піддослідних тварин через 21 день після змодельованого опікового ушкодження при системному застосуванні NaCl можна охарактеризувати як адаптаційні та компенсаторні процеси в стінці фолікула, судинному та стромальному компонентах. Більшість фолікул щитовидної залози були круглої або овальної форми, деякі з них були перерозтягнутими і заповнені щільним колоїдом. Клітини фолікулярного епітелію були сплющені, містили пікнотичні ядра, оточені тонким шаром цитоплазми. У деяких регіонах спостерігали десквамацію тироцитів у просвіті фолікула. Електронно-мікроскопічне дослідження залози через 21 день після змодельованого опіку підтвердило зміни, які раніше були встановлені аналізом гістологічних препаратів. Сплощені тироцити містили осміофільні ядра з гетерохроматином. Їх цитоплазма була електронно прозорою і містила органели з ознаками руйнування. Мікроворсинки спостерігали на апікальній області тироцитів. Поряд із фолікулами зі зміненою гістологічною структурою (що свідчить про дисбаланс у синтезі та виведенні гормонів) спостерігалися фолікули з інтактною стінкою та неушкодженим кровопостачанням. Ультраструктура останнього свідчить про активні білково-синтезуючі та білково-секреторні процеси.

Ключові слова: опікова травма, щитовидна залоза, світлова мікроскопія, електронна мікроскопія.

The work is fragments of the research project "Peculiarities in micro-/ultramicroscopic structure and histochemical properties of body tissues during the development of compensatory-adaptive reactions", state registration No 0121U108204.

It is widely known that severe burn immediately subjects the human body to extreme stress which leads to a series of immunological, neuroendocrine and metabolic reactions [11, 13]. Although the initial response to severe burn is typically characterized by hypometabolism, this state quickly (72–96 hrs) changes to hypermetabolism [4, 7]. The post-burn hypermetabolism is believed to be caused by the